

Quantum Transport of a 1D Degenerate Bose Gas in a Lattice: The Role of Different Quantum Fluctuations

B. Xiong¹, T. Yang¹, S. Chen², and Keith A. Benedict¹

¹ *Midlands Ultracold Atom Research Center, School of Astronomy and Physics,
University of Nottingham, Nottingham NG7 2RD, United Kingdom*

² *Department of Physics, Wuhan University, Wuhan 430072, People's Republic of China*
(Dated: February 10, 2010)

We study the effect of quantum fluctuations on the quantum dynamics of 1D Bose gases in an optical lattice using different basis expansions and the Truncated Wigner approximation. The initial spatial distributions of the quantum fluctuations are very different when using a limited number of plane-wave (PW), single-harmonic-oscillator (SHO), and the self-consistently determined Bogoliubov (SCB) modes. The transport properties of the Bose gas, characterized by the phase coherence and number fluctuations, in the PW basis are distinct from those gained using SHO and the SCB basis. The calculations using the SCB modes predict greater phase decoherence and stronger number fluctuations than the other choices. Furthermore, we observe that the use of PW modes overestimates the extent to which atoms are expelled from the core of the cloud, while the use of the other modes only breaks the cloud structure slightly which is in agreement with the experimental observations [1].

PACS numbers: 03.75.Dg, 03.75.Lm, 39.20.+q

I. INTRODUCTION

The influence of quantum fluctuations on ultracold atomic gases have attracted much attention in recent years. In periodic optical lattice (OL), bosons can undergo a superfluid-to-Mott-insulator (SF-MI) quantum phase transition because of the loss of phase coherence induced by quantum fluctuations if the tunneling became small enough [2, 3]. Theoretical work has shown that for a system where condensate oscillations are overdamped in a 1D OL, there is a smooth crossover between the classical transition and the SF-MI phase transition at zero displacement as the quantum fluctuations increase [4]. In addition, the damped motion of a 1D Bose gas in a shallow OL has been demonstrated in experiment [1]. Many theoretical studies have indicated that quantum fluctuations can result in strong dissipation which leads to dynamical instabilities [4–7].

The truncated Wigner approximation (TWA) developed in the field of quantum optics provides an attractive framework to explore the effect of the quantum fluctuations on the dynamics of Bose-Einstein condensates (BECs) [5, 8–15]. The fundamental idea of this method is to expand quantum field operators in the equation of motion for a Bose gas system in the Wigner representation and, by the neglect of the third-order functional derivatives in the equation, derive a generalized Fokker-Planck equation (FPE) for a classical field problem. This results in a stochastic differential equation for the classical field $\Psi(\mathbf{x}, t)$, which coincides with the mean field Gross-Pitaevskii equation [5, 13]

$$i\hbar\partial_t\Psi(\mathbf{x}, t) = H_0\Psi(\mathbf{x}, t) + g|\Psi(\mathbf{x}, t)|^2\Psi(\mathbf{x}, t). \quad (1)$$

where $H_0 = -\hbar^2\partial_{\mathbf{x}}^2/2m + V(\mathbf{x}, t)$. The quantum-mechanical vacuum fluctuations are simulated by adding appropriate classical fluctuations in addition to the co-

herent field of the *initial* state of the BEC. Specifically, the quantum fluctuations are introduced in the *initial condition* for the classical field $\Psi(\mathbf{x}, t)$, by the sum $\Psi(\mathbf{x}, t=0) = \psi(\mathbf{x}) + \xi(\mathbf{x})$, where $\psi(\mathbf{x})$ and $\xi(\mathbf{x})$ are the real and virtual particle fields, respectively. The real particle field is the suitably chosen initial form for the classical field. The field of virtual particles is decomposed into a basis set $\xi(\mathbf{x}) = \sum_{j=1}^M \xi_j \phi_j(\mathbf{x})$, where $\phi(\mathbf{x})$ is the mode function and the amplitude in each mode is Gaussian with the properties, $\langle \xi_i^* \xi_j \rangle = \frac{1}{2} \delta_{i,j}$, $\langle \xi_i \xi_j \rangle = 0$. Expectation values are evaluated by averaging over an ensemble of mean-field calculations with random initial conditions. It implies that the stochastic element only appears in the initial conditions.

Recently, the TWA has predicted quantum turbulence in the collisions of two condensates [13] and explained the disruption of reflecting BECs from Silicon surface [14]. In these theoretical works, zero-temperature quantum fluctuations are modeled by adding, on average, a half quantum into each low-energy PW mode. Therefore, the quantum fluctuations added into the nonuniform condensates are spatially uniform, suggesting that the distribution of quantum fluctuations does not depend on the geometry of atom clouds and interatomic interaction. However, if the quantum is added into a *limited* number of *distinct* modes in the treatment of TWA, the corresponding initial wave functions of the classical field are different. In addition, the wave vector is not a good quantum number for a harmonically trapped condensates [16, 17], except for a 1D, very elongated, condensate [15]. In the frame of Bogoliubov theory, the geometry and interatomic interaction of condensate clouds have a critical effect on the distribution of quantum fluctuations. The TWA with Bogoliubov theory in the treatment of quantum fluctuations has shown reasonable results in the nonadiabatic dynamics and dissipative dynamics of quan-

tum transport in optical lattices [5]. Provided a complete set of modes are used, the choice of basis would be immaterial from the point of view of the validity of the method, and the computationally simplest basis is preferred. However, the method requires a cut-off to be imposed on the mode expansion which distinguishes “low energy” modes, that properly contribute to fluctuations in the condensate, from “high energy” modes that do not. The use of a truncated basis means that it is no longer clear that all basis sets are equivalent.

In this paper, we consider three choices of basis set: PW modes, SHO eigenstates, and SCB modes, to construct the quantum fluctuations with different initial states in TWA. Our purpose is to examine the effect of these different quantum fluctuations on the dynamics of a strongly confined Bose gas, oscillating in a very shallow 1D OL. While we find, as shown in Ref. [1, 5], that the inclusion of quantum fluctuations does indeed lead to damping of the dipole oscillations of the center-of-mass (c.m.) of the condensate, we note that the choice of restricted basis set has a qualitative effect on the evolution of the spatial density distribution of the cloud, the loss of phase coherence, and the number fluctuations. These qualitative differences originate from the initial distribution of the quantum fluctuations. In the PW basis, some atoms are kicked out from the core part of atomic cloud, resulting in a density distribution with a long tail. This does not agree completely with the experimental observations [1], where the cloud in damped transport has a similar width to the undamped case. However, there is only slight disruption of the cloud structure in the other two cases. Moreover, by using the same number of modes, Bogoliubov theory offers stronger damping of the c.m. trajectory than others.

The remainder of this paper is organized as follows. In section II we summarize the truncated Wigner scheme and our specific implementation of it for the three choices of basis states. In section III we describe the results of the simulations and highlight the differences in dynamical evolution between the different choices of basis set. In section IV we summarize these and draw more general conclusions about such simulations.

II. THEORETIC MODEL AND NUMERICAL METHODS

Our theoretical model is based on the recent experiment [1], where damped dipole oscillations of a 1D Bose gas in a shallow OL were observed. In this experiment, the 1D Bose gas is formed in an array of independent “tubes”, produced by applying a strong transverse 2D OL potential to confine a trapped 3D atomic condensate. Then the tubes are corrugated adiabatically by using a very shallow 1D lattice along the axial direction. The dipole oscillations of atoms along the weak axial lattice were excited by suddenly displacing the harmonic trap and the c.m. velocity was imaged.

The transport dynamics of the system based on the above model is studied by using the TWA method. In a very shallow OL, the dynamics of the system can be described by the 1D GPE (Eq.(1)) with the coupling constant $g = g_{1D} = 2\hbar\omega_{\perp}a$. We use the s-wave scattering length $a = 5\text{nm}$ for ^{87}Rb and $\omega_{\perp} = 2\pi \times 38\text{kHz}$, originated from the quadratic expansion of the strong transverse 2D OL around the local minima [1, 5]. The potential energy profile of the 1D optical lattice in the axial direction is characterized by $V_{OL}(x, t) = A(t)\sin^2(\pi x/d)$, where $d = 405\text{nm}$ with respect to the laser wavelength $\lambda = 810\text{nm}$. The amplitude $A(t)$ is assumed to zero initially, ramped gradually up to $E_r/2$ in 2.65ms as $\exp(kt) - 1$ where $E_r = \hbar^2/2m\lambda^2$ is the photon recoil energy, and k is a constant determined from the ramping time. Thereafter $A(t)$ remains unchanged and the total confining potential is $V(x, t) = V_{OL}(x, t) + m\omega^2x^2/2$, where $\omega = 2\pi \times 60\text{Hz}$ is angular frequency of the harmonic trap. At time $t = 2.65\text{ms}$, we abruptly displace the harmonic trap through a distance $\Delta x = 3\mu\text{m}$, accelerating the Bose gas in the OL.

We consider the dynamics of the system for distinct initial states by introducing different mode functions $\phi(x)$ in the field of virtual particles $\xi(x)$. In case I, the condensate is assumed to propagate in the OL in absence of quantum fluctuations, i.e. $\xi^*(x)\xi(x) = 0$. In case II, the initial amplitude of virtual particle field, $\langle \xi_i^* \xi_i \rangle$, is 1/2 and $\phi_i(x) = \exp(ik_i x)/\sqrt{L}$, where k_i is the wave vector in the i -th mode and L is the length of system [13, 14]. In case III, the mode function, $\phi_n(x)$, is the eigenfunction of the SHO, resulting in the quantum fluctuations in a limited number of energy modes. In case IV, we use Bogoliubov approximation

$$\hat{\Psi}(x, t) = \psi_0(x)\hat{a}_0 + \sum_{j>0} [u_j(x)\hat{a}_j(t) - v_j^*(x)\hat{a}_j^\dagger(t)], \quad (2)$$

where $\psi_0(x)$ is the ground state solution of the GPE, the number of atoms $\langle \hat{a}_0^\dagger \hat{a}_0 \rangle = N_0$, and $\hat{a}_j^\dagger, \hat{a}_j$ are quasiparticle field operators. Then we diagonalize the Bogoliubov equations,

$$\begin{aligned} \mathcal{L}u_j(x) - g_{1D}N_0\psi_0^2(x)v_j(x) &= E_j u_j(x) \\ \mathcal{L}v_j(x) - g_{1D}N_0\psi_0^{*2}(x)u_j(x) &= -E_j v_j(x), \end{aligned} \quad (3)$$

in the subspace *orthogonal* to the condensate space to obtain the quasiparticle amplitudes, $u_j(x)$, $v_j(x)$ and excitation energy E_j . In Eqs.(3), $\mathcal{L} = -\hbar^2\partial_x^2/2m + U_{eff}(x)$ with

$$U_{eff}(x) = V(x) + 2g_{1D}N_0|\psi_0(x)|^2 - \mu \quad (4)$$

the effective potential of quasiparticles [18]. The initial state of $\Psi(x)$ can be obtained using Eq.(2) and $\hat{a}_j, \hat{a}_j^\dagger$ ($j > 0$) are replaced by the random variables (α_j, α_j^*) , obtained by sampling the corresponding Wigner distribution of the quasiparticles in a thermal bath [19],

$$W(\alpha_j, \alpha_j^*) = \frac{2}{\pi} \tanh(\eta_j) \exp[-2|\alpha_j|^2 \tanh(\eta_j)], \quad (5)$$




FIG. 1: The spatial distribution of density fluctuations, attributed respectively by adding a half quantum into each PW (a), SHO (b), and SCB mode (c). The effective potential U_{eff} defined by Eq.(4) (d).

where $\eta_j = E_j/2k_B T$. At temperature $T = 0$, the quantum fluctuations are described by $\xi(x) = \sum_j u_j(x)\alpha_j - v_j(x)\alpha_j^*$ in the SCB modes based on Eq.(2) and Eq.(5). In this paper, we consider the total atom number $N_0 = 100$, the number of modes, $M < 80$, and always restrict our discussions at zero temperature where the Bogoliubov approximation can be justified. Two groundstates with optical lattice amplitudes $A = 0$ and $E_r/2$ are calculated numerically by evolving the GPE in imaginary time [20].

The plane waves are the simplest computationally as the field of virtual particles $\xi(x)$ is obtained using the Fast Fourier transform method. The SHO basis has the advantage that the support of each mode is weighted towards the region in which the cold atoms are likely to be found and, while the construction of $\xi(x)$ is more complicated, the modes are at least pre-defined and are relatively straightforward to generate iteratively. The SCB modes should be good approximations to the quasi-particle excitations of the system, at least for weak interactions, and are well adapted to the specifics of the potential in which the cloud moves. The calculation of $\xi(x)$ is more computationally demanding because the form of the modes has to be determined by diagonalization of a large matrix whose elements depend on the form chosen for the initial order parameter field, $\psi_0(x)$.

To determine the validity and precision of the results, different numerical methods are applied in our simulation. In case I, we obtain dynamic results by using the Crank-Nicolson method to solve the time-dependent Gross-Pitaevskii equation. In case II, a distinctive numerical method, RK4IP-P [13, 21], is utilized. Both methods are used in case III-IV to examine their equivalence. Solutions of the c.m. trajectory, phase coherence and number fluctuation in TWA, are calculated numerically using different numbers of realizations. We find that some results can be significantly different when the number of realizations is small, i.e. less than 50. Therefore, we show all results based on 200 realizations for




FIG. 2: (Color online) The time evolution of the c.m. motion of the Bose gas in cases I (solid black line), II (solid green line), III (dashed red line), and IV (solid blue line). The fitting of the c.m. trajectories in cases III and IV (dotted black line in (b) and (d)) from Eq.(8). The corresponding parameters: $B = 0.22\mu\text{m}$, $0.1\mu\text{m}$ and $\gamma = 22.89, 10.44$ for cases III and IV, respectively, while $\Omega = 2\pi \times 60$ and $A = 3.0\mu\text{m}$ for both cases.

numerical consistency.

III. RESULTS AND DISCUSSION

In Fig.1, we show the initial spatial distributions of the quantum fluctuations $\langle \xi^*(x)\xi(x) \rangle$ with respect to three sets of basis modes. In the PW modes, $\langle \xi^*(x)\xi(x) \rangle = \frac{1}{2L} \sum_{i,j=1}^M \langle \xi_i^* \xi_j \rangle \exp[i(k_i - k_j)x]$, indicating a uniform spatial distribution of the quantum fluctuations, while the *limited* number of low-energy SHO modes are dominated by the Gaussian function, which induces a localized spatial distribution of the quantum fluctuations. In contrast, the quantum fluctuations in the SCB modes give a spatial distribution with a double peak beside the atom cloud, which coincides with the effective potential of the quasi-particles Eq.(4) [18], as shown in Fig.1(d).

Qualitative insight into the properties of the quantum dynamics can be gained by using classical dissipative dynamics to compare with our quantum simulations. We model the c.m. motion as a damped harmonic oscillator [1, 5]

$$\ddot{X}_{c.m.} + 2\gamma\dot{X}_{c.m.} + \frac{k}{m}X_{c.m.} = 0 \quad (6)$$

where the c.m. displacement is defined by

$$X_{c.m.} \equiv \langle x \rangle(t) = \frac{\int_{-\infty}^{\infty} x |\Psi(x, t)|^2 dx}{\int_{-\infty}^{\infty} |\Psi(x, t)|^2 dx}. \quad (7)$$




FIG. 3: (Color online) Phase coherence C_3 in different cases (a), (b), and C_j in case III (curves from top C_3 , C_5 , C_8 , C_{10}) (c).

In underdamped case, Eq.(6) have a solution

$$X_{c.m.} = -e^{-\gamma(t-t_0)} [A \cos \Omega(t-t_0) + B \sin \Omega(t-t_0)] \quad (8)$$

with $B = \gamma A / \Omega$, $\Omega = \sqrt{k/m - \gamma^2}$, and t_0 determined by the initial phase.

In Fig.2, we show simulation results of the c.m. trajectories for case I-IV in comparison with Eq.(8). In case I, there is undamped motion of the c.m. while the visibly damped trajectories occur in presence of the quantum fluctuations in the other cases. The damped trajectories $\langle x \rangle(t)$ cannot be fitted by Eq.(8), especially at long times, highlighting the complexity of the effect of the quantum fluctuations on the transport of the system. Therefore, the chosen fitting parameters satisfy the numerical calculations well in the first period but hardly characterize the subsequent behavior of the damped motion. To compare the time-dependent damping rate in different cases, we introduce $\lambda \equiv \ln(D_0/D_1)$ [4], where D_0 and D_1 are the c.m. positions at $t = 10.9\text{ms}$ and 27.6ms (Fig.2(c)). Our calculations for cases III and IV (Fig.2(c)) give $\lambda = 0.21$ and 0.26 respectively, therefore indicating that, with the same number of basis states, the quantum fluctuations in SCB modes inhibit more strongly the motion of the Bose gas than in other modes. There are slight differences between the c.m. trajectories in cases II and III, as shown in Fig.2(a).




FIG. 4: (Color online) Number fluctuation Δn_0 in different cases (a), (b), and Δn_j in case III (curves from top Δn_0 , Δn_{10} , Δn_{20} , Δn_{40}) (c).

The damped motion of the c.m. suggests that there might be a loss of phase coherence, and number fluctuations in the system. In order to avoid the complications arising from the symmetrically ordered multimode field Ψ [5], we define the ground state operators b_j for each individual lattice site j :

$$b_j(t) = \int_{j^{th} well} dx \Psi_0^*(x) \Psi(x, t), \quad (9)$$

where $\Psi(x, t)$ is the stochastic field, determined by Eq.(1) and $\Psi_0(x)$ is the groundstate wave function in the combined harmonic trap and OL. The normalized phase coherence between the central well and its j th neighbor and the atom number fluctuations in the j th site are separately set as $C_j = |\langle \hat{b}_0^\dagger \hat{b}_j \rangle| / \sqrt{n_0 n_j}$ and $\Delta n_j = [\langle (\hat{b}_j^\dagger \hat{b}_j)^2 \rangle - \langle \hat{b}_j^\dagger \hat{b}_j \rangle^2]^{1/2}$.

In Fig.3 and Fig.4, we show respectively the configurations of the dynamic phase coherence and number fluctuations for cases II-IV ((a) and (b)) and on different spatial sites for case III ((c)). In case III, there is an increase of the phase coherence C_3 at $t < 4.8\text{ms}$ and then its loss at $t > 4.8\text{ms}$ in the first period of the c.m. oscillation, while the number fluctuations Δn_3 decrease and grow correspondingly, displaying the opposite dynamic behavior between the phase coherence and number fluctuations at nearby sites. This behavior demon-




FIG. 5: Density distribution at $t = 8.65\text{ms}$ in cases II (a), III (b) and IV (c). $l_b = \sqrt{\hbar/m\omega}$ is the length unit.

strates that the quantum fluctuations tend to suppress the phase coherence. Moreover, the phase coherence C_j and number fluctuation Δn_j are correlated essentially to the damped trajectory of the c.m.. At $t = 2.65\text{ms}$, the c.m. in Fig.2 is far from the center of the trap potential, and thus C_3 and Δn_0 , as shown in Fig.3(c) and Fig.4(c), identify the properties in the outer region of the Bose gas where the number fluctuation is large and the phase coherence is weak. Then the Bose gas moves toward the center of the trap, and Δn_0 decreases gradually while C_3 increases.

However, qualitative differences are particularly marked at short times, as depicted in Fig.3(a) and Fig.4(a), indicating at least a marked dependence of the “equilibration time” for such simulations on the choice of basis set. This is most marked for the PW basis in which the short-time oscillations in the fluctuation statistics are quite different to the others and only approach them at times rather longer than the period of the dipole oscillations. Fig.3 (b) and Fig.4 (b) show that the short-time variation of C_3 and Δn_0 based on the SCB modes is more complicated than the other two modes. This might originate from the greater complexity of the initial distribution of the quantum fluctuations in the SCB modes. Comparing the phase coherence for cases III and IV, shown in Fig.3(b), we find that the time average of C_3 from 2.65ms to 42.65ms for case IV is 0.63, 5.21% smaller than that for case III (0.66). The corresponding mean value of the number fluctuation in the time interval for case IV is 3.95, a little higher than 3.88 for case III. These indicate that with the same number of basis

modes, the quantum fluctuations in the SCB modes cause stronger phase decoherence and number fluctuation than those in the PW and SHO modes.

Interestingly, we also find that, during the transport process, some atoms are kicked out from the core region of the atom cloud in case II (Fig.5(a)) while the quantum fluctuations only break the inner configuration of the condensate slightly in cases III (Fig.5(b)) and IV (Fig.5(c)). In the experiment [1], they did not observe a significant difference in the time of flight width between atoms that undergo damped harmonic motion and those that are unexcited but held for an equal time.

Of course, we can strictly only say that the three basis sets produce different results, but it is natural to assume that the simulations using the SCB modes have more information about the specific physics of the system built into them and should be the most reliable. We then note that for some properties, the gain in using the SCB modes compared to the SHO modes is small given the considerable increase in computational complexity. However we note that in the simulation that uses PW modes for trapped BECs, a long equilibration time should be allowed before any non-trivial dynamical processes are allowed to occur in the simulation to avoid spurious transient effects.

IV. CONCLUSION

In conclusion, we have studied the effect of different choices of basis set for the inclusion of quantum fluctuations on a 1D Bose gas in an OL using the TWA. Specifically we have used PW, SHO and the SCB modes to decompose the quantum fluctuations and examine the dynamics of the system. The difference in the predictions of the phase coherence, number fluctuations and density variations at short time indicates that the choice of the basis set has a substantial influence on the qualitative features of the transport. The use of the SCB modes, which gives greater phase decoherence and stronger number fluctuations on the damping dynamics than the other two choices, incorporates more of the underlying physics of the problem and presumably leads to the most reliable results. For the trapped condensate, the SHO basis give results that are qualitatively similar to those from the SCB modes, at rather less computational effort. It’s therefore clear that it is important to select the basis set that best suits the particular system to be studied.

ACKNOWLEDGMENTS

We thank R. G. Scott for fruitful discussion. We also acknowledge support by the EPSRC.

-
- [1] C. D. Fertig, K. M. O'Hara, J. H. Huckans, S. L. Rolston, W. D. Phillips, and J. V. Porto, Phys. Rev. Lett. **94**, 120403(2005).
 - [2] D. Jaksch, et al., Phys. Rev. Lett. **81**, 3108 (1998).
 - [3] M. Greiner, et al., Nature (London) **415**, 39(2002).
 - [4] A. Polkovnikov and D.-W. Wang, Phys. Rev. Lett. **93**, 070401 (2004).
 - [5] J. Ruostekoski and L. Isella, Phys. Rev. Lett. **95**, 110403(2005); L. Isella and J. Ruostekoski, Phys. Rev. A **72**, 011601(2005); L. Isella and J. Ruostekoski, Phys. Rev. A **74**, 063625(2006).
 - [6] J. Gea-Banacloche et al., cond-mat/0410677.
 - [7] E. Altman et al., Phys. Rev. Lett. **95**, 020402 (2005).
 - [8] M. J. Steel, M. K. Olsen, L. I. Plimak, P. D. Drummond, S. M. Tan, M. J. Collett, D. F. Walls, and R. Graham, Phys. Rev. A, **58**, 4824(1998).
 - [9] H. T. C. Stoof and M. J. Bijlsma, J. Low Temp. Physics **124**, 431(2001).
 - [10] M. J. Davis, S. A. Morgan, and K. Burnett, Phys. Rev. Lett. **87**, 160402(2001).
 - [11] A. Sinatra, C. Lobo, and Y. Castin, Phys. Rev. Lett. **87**, 210404(2001).
 - [12] K. Góral, M. Gajda, and K. Rzażewski, Phys. Rev. A **66**, 051602(R)(2002).
 - [13] A. A. Norrie, R. J. Ballagh, and C. W. Gardiner, Phys. Rev. Lett. **94**, 040401(2005); Phys. Rev. A **73**, 043617(2006).
 - [14] R. G. Scott, D. A. W. Hutchinson, and C. W. Gardiner, Phys. Rev. A **74**, 053605(2006); Laser Physics **17**, 527(2007).
 - [15] R. G. Scott, T. E. Judd, and T. M. Fromhold, Phys. Rev. Lett. **100**, 100402(2008); R. G. Scott, D. A. Hutchinson, T. E. Judd, and T. M. Fromhold, Phys. Rev. A, **79**, 063624(2009).
 - [16] R. Ozeri, N. Katz, J. Steinhauer, and N. Davidson, Rev. Mod. Phys. **77**, 187(2005).
 - [17] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Oxford: Clarendon) (2003).
 - [18] A. Csordás, R. Graham, and P. Szépfalusy, Phys. Rev. A **57**, 4669(1998).
 - [19] C. W. Gardiner and P. Zoller, *Quantum Noise* (Springer, Berlin, 1999).
 - [20] M. L. Chiofalo, S. Succi, and M. D. Tosi, Phys. Rev. A **62**, 7438(2000).
 - [21] *A Classical Field Treatment of Colliding Bose-Einstein Condensates*, A. A. Norrie, A thesis submitted for the degree of Doctor of Philosophy at the University of Otago, Dunedin, New Zealand, June 2005.